



#### WHAT IS IT?

Sterile cosmeceutical solution with tranexamic acid, kojic acid, ascorbic acid (vitamin C), nicotinamide, rutine and high, medium and low molecular weight hyaluronic acid, with moisturizing and depigmenting action.

The formulation of BSK 14 has been reinforced with Kojic acid and vitamin C for greater antioxidant action, reducing Hyperpigmentation significantly.

Formulated with hyaluronic acid, with a moisturizing activity, an increase in elasticity and an activation of collagen and elastin production due to its low molecular weight fraction.

With vitamins that reinforce and restructure the dermal layer have been used for rapid recovery after treatment.

To prolong the effects of the treatment, it is recommended to use Serum BSK76 and Depigmenting Cream BSK70.



#### **PRESENTATION**

Vials 5ml / 0.175 Fl. Oz. | 1 Box of 5 uds

### **USE**

Cosmetic product. Topical use. Facial and hands product.

## **INGREDIENTS**

Aqua Tranexamic Acid Ascorbic Acid Sodium Hyaluronate Kojic Acid Niacinamide Glutathione Troxerutin

## **PRECAUTIONS**

- Preserve between 5°C y 25°C
- Keep out of children's reach
- Avoid contact with eyes and mucous membranes
- Suspend its use if there is allergy, sensitivity or hipersensitivity to the product



#### **HOW TO USE**

How to apply?

Apply on the area to be treated, giving a light massage until product complete absorption.

Suitable for cosmetic treatments and virtual mesotherapy.

External use. Topical use.


Facial and hands product

		~	•		_			
ப	roi	-	10r		Irc	\Ot	m	nt
г	I OI	fess	IUI	ıuı	116	7UL	ו ווכ	71 IL

- 1. Using the transdermal therapy technique chosen by the professional, apply 3-5 ml of the solution (BSK14)
- 2. Repeat the application after three weeks
- 3. For effective treatment between the first and second professional sessions, tranexpro BSK76, BSK70 and sunscreen products should be applied daily at home
- 4. Repeat as required to achieve treatment objectives every 6-12 months

	Н	lome	Trea	tment
--	---	------	------	-------

- 1. Clean your face and neck in the morning and at night
- 2. Apply the Face Tonic
- 3. After cleaning your face apply the Serum Tranexpro on the area to be treated, giving a light massage until product complete absorption
- 4. After the serum, apply Cromabright cream giving a light massage until product complete absorption
- 5. In the morning, it's recommended, apply Solar protector





## **STUDIES**

There are currently different studies that support the efficacy of topical use of 5% tranexamic acid to reduce hyperpigmentation due to long sun exposure, hormonal changes, what is commonly known as melasma or ect.

Studies have been done comparing the efficacy of this active against hydroquinone, a product that cannot be used in cosmetic products.

# Art.1 **Tranexamic Acid for Melasma: Evaluating the Various Formulations.**Wang JV, Jhawar N, Saedi N. - J Clin Aesthet Dermatol. 2019;12(8): E73-E74.

## Art. 2 **Topical tranexamic acid as a promising treatment for melasma.**Ebrahimi B, Naeini FF - J Res Med Sci. 2014;19(8): 753-757.

# Art. 3 A Randomized Controlled Study Comparing the Efficacy of Topical 5% Tranexamic Acid Solution versus 3% Hydroquinone Cream in Melasma.

Janney MS, Subramaniyan R, Dabas R, Lal S, Das NM, Godara SK. J Cutan Aesthet Surg. 2019;12(1):63–67. doi:10.4103/JCAS.JCAS\_40\_18




Gráfico 3 - Patient self-assessment of improvement at week 12. Patients were asked to grade their overall improvement as 1 = 75% lightening (excellent), 2 = 51-75% (good), 3 = 26-50% (fair), and 4 = 0-25% lightening (poor)
Font. Journal of Research in Medical Sciences




Gráfico 2 - Physician assessment of improvement at week 12 based on the patients photos. Physician were asked to grade their overall improvement as 1 = :75% lightening (excellent), 2 = 51-75% (good), 3 = 26-50% (fair), and 4 = 0-25% lightening (poor) Font. Journal of Research in Medical Sciences

## **STUDIES**




Gráfico 1 - Changes of Melasma Area and Severity Index score of two groups Font: Journal of Research in Medical Sciences




Gráfico 4 - Melasma area severity index (MASI) scores within and between tranexamic acid (TA) and hydroquinone (HQ) groups at baseline, and after 4, 8, and 12 weeks

Font: Journal of Cutaneous and Aesthetic Surgery



## PRODUCT IMAGE



Preserve between 5°C y 25°C. Keep out of children's reach. Avoid contact with eyes and mucous membranes. Suspend its use if there is allergy, sensitivity or hipersensitivity to the product.

Cosmetic product. Topical use. External use.



## **ADDITIONAL PRODUCTS**



**BSK76** TRANEXPRO

> Serum 30ml



**BSK70**TRANEXPRO
Depigmentation

Cream 50ml



Recommended volume (ml) per area

# Recommended depth per area

